Diagnostik der TTP

Ralph Wendt
Abteilung Nephrologie
Klinik für Infektiologie/Tropenmedizin, Nephrologie und Rheumatologie

ISTH guidelines for the diagnosis of thrombotic thrombocytopenic purpura

X. Long Zheng¹ | Sara K. Vesely² | Spero R. Cataland³ | Paul Coppo⁴ | Brian Geldziler⁵ | Alfonso Iorio^{6,7} | Masanori Matsumoto⁸ | Reem A. Mustafa⁹ | Menaka Pai⁷ | Gail Rock¹⁰ | Lene Russell¹¹ | Rawan Tarawneh¹² | Julie Valdes¹³ | Flora Peyvandi^{14,15}

COI

lecture fees / advisory board honaria / travel grants:

Fresenius Medical Care, Vifor Fresenius Pharma, Amgen, Shire, Novartis, Alexion, Otsuka, Cellpharm, **Ablynx**, Daiichi Sankyo, **Sanofi**, AstraZeneca

Agenda

- 1) Thrombotische Mikroangiopathie – What's that?
- 2) Diagnose der TTP
- 3) Was sagen die ISTH Leitlinien?
- 4) Zusammenfassung

Thrombotische Mikroangiopathie

Fragmentozyten

Fragmentozyten als klassische Helmform bei TTP

Fragmentierte Erythrozyten (Schizozysten) bei TTP

Fragmentozyten bei HUS

Akanthozyten-ähnliche Formen bei HUS

Differentialdiagnose der TMA

Differential Diagnosis for Thrombotic Microangiopathies (TMAs)

ADAMTS13

A Disintegrin And Metalloprotease with a ThromboSpondin type 1 motif, member 13

Klinikum | **St.GEORG**

INSELSPITAL

UNIVERSITÄTSSPITAL BERN
HOPITAL UNIVERSITAIRE DE BERNE
BERN UNIVERSITY HOSPITAL

Die Entdeckung der ADAMTS13: 1998 in Bern

[Baillieres Clin Haematol.](#) 1998 Jun;11(2):509-14.

Deficiency of von Willebrand factor-cleaving protease in familial and acquired thrombotic thrombocytopenic purpura.

Furlan M¹, Lämmle B.

⊕ Author information

Abstract

Excessive intravascular platelet agglutination in patients with thrombotic thrombocytopenic purpura (TTP) appears to be associated with excessive release from endothelial cells of unusually large von Willebrand factor (vWF) multimers and/or impaired degradation of these multimers by a 'depolymerase' cleaving vWF to smaller, non-agglutinating molecular forms. We studied the activity of a recently described vWF-cleaving protease in four patients, including two brothers, with chronic relapsing TTP. All four patients had lacking or strongly reduced vWF-cleaving protease activity. In another patient with chronic relapsing TTP, the protease deficiency was due to the presence in the patient plasma of an inhibitor that was found to be an IgG. We conclude that constitutional as well as acquired deficiency of vWF-cleaving protease may predispose to clinical manifestation of TTP.

PMID: 10097823

The New England Journal of Medicine

VON WILLEBRAND FACTOR-CLEAVING PROTEASE IN THROMBOTIC THROMBOCYTOPENIC PURPURA AND THE HEMOLYTIC-UREMIC SYNDROME

MIHA FURLAN, PH.D., RODOLFO ROBLES, MIRIAM GALBUSERA, Sc.D., GIUSEPPE REMUZZI, M.D., PAUL A. KYRLE, M.D., BRIGITTE BRENNER, MANUELA KRAUSE, M.D., INGE SCHARRER, M.D., VOLKER AUMANN, M.D., UWE MITTLER, M.D., MAX SOLENTHALER, M.D., AND BERNHARD LÄMMLE, M.D.

Prof. Dr. Bernhard Lämmle

Emeritus Professor und Direktor Universitätsklinik für
Hämatologie, Bern
Senior Professor am CTH der Universitätsmedizin Mainz

TTP: Prä-Test-Risikoabschätzung und Diagnose

- Plasmic (oder auch French-) Score
- ADAMTS13 Schnelltest
- ADAMTS13 Aktivität im (spezialisierten) Labor (+ Inhibitor/Antikörper)

Algorithmus TMA-Abklärung

Unterscheidung TTP und aHUS

Hohe Thrombozytenzahlen ($>30,000 \text{ mm}^3$) und S-Kreatininwerte ($>150\text{--}200 \mu\text{mol/l}$) schließen die Diagnose einer schweren ADAMTS13-Defizienz (TTP) nahezu aus

Patientencharakteristika*	TTP	aHUS	p-Wert
	ADAMTS13 defizient n=160 (Standardabweichung)	ADAMTS13 messbar n=54 (Standardabweichung)	
Thrombozytenzahl, $\times 10^9/l$	17,4 (14,2)	66,6 (49,3)	<0,0001
Kreatininlevel, $\mu\text{mol/l}$	114 (68,4)	454 (326)	<0,0001

TTP Plasmic score

Parameters	French Score	PLASMIC Score
Platelet count	<30 × 10 ⁹ /L (+1)	<30 × 10 ⁹ /L (+1)
Serum creatinine level	<2.26 mg/dL (+1)	<2.0 mg/dL (+1)
Hemolysis		
Indirect bilirubin >2 mg/dL or reticulocyte count >2.5% or undetectable haptoglobin	a	+1
No active cancer in previous year	a	+1
No history of solid organ or SCT	a	+1
INR < 1.5	a	+1
MCV < 90 fL	NA	+1
Likelihood of severe deficiency of ADAMTS13 activity (<10%)	0: 2%	0-4: 0%-4%
	1: 70%	6: 5%-24%
	2: 94%	6-7: 62%-82%

Variables	Points*
Platelet count <30 x 10 ⁹ per L	1
Hemolysis variable†	1
No active cancer	1
No history of solid-organ or stem-cell transplant	1
MCV <90 fL	1
INR <1.5	1
Creatinine <2.0 mg/dL	1
INR: International normalized ratio. MCV: Mean corpuscular volume. †Reticulocyte count >2.5%, or haptoglobin undetectable, or indirect bilirubin >2.0 mg/dL.	

TTP Plasmic score Validierungen

	Derivation cohort (n=200)	Internal validation cohort (n=150)	External validation cohort (n=146)
0-4	0/84 (0%)	0/89 (0%)	2/47 (4%)
5	2/44 (5%)	3/32 (9%)	6/25 (24%)
6 or 7	58/72 (81%)	18/29 (62%)	61/74 (82%)

Data are number of individuals with ADAMTS13 activity of 10% or less/total number of individuals with that score (%).

Score	Risk Category	Risk of severe ADAMTS13 deficiency ($\leq 10\%$)
0-4	Low	4.3%
5-6	Intermediate	56.8%
7	High	96.2%

Schnelltest versus ELISA

TABLE 4 Misclassified results at ADAMTS-13 activity cutoff of 0.1 IU/mL

Group	n	Subgroup (n)	Quantitative ADAMTS-13 activity (IU/mL)	TECHNOSCREEN® (IU/mL)
1	12	5	0.00/0.00/0.04/0.07/0.09	0.1
		7	0.10/0.11/0.12/0.14/0.15/0.17/0.18	0
2	9	-	0.33/0.38/0.39/0.40/0.42/0.55/0.71/0.85/0.86	0
3	1	-	0.00	0.4

ADAMTS13 Schnelltest

Normal sample
ADAMTS-13 activity: 0.95 IU/mL by ELISA
0.8 IU/mL by screen

TTP sample
ADAMTS-13 activity: 0.05 IU/mL by ELISA
0.0 IU/mL by screen

Site	n	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV % (95% CI)	NPV % (95% CI)	Accuracy % (95% CI)
All	220	88.7 (77.4-94.7)	90.4 (85.0-94.0)	74.6 (62.7-83.7)	96.2 (91.9-98.2)	90.0 (85.3-93.6)

ISTH Leitlinie verwendet 3 Szenarien:

1) Szenario A: ADAMTS13 Aktivität *jederzeit* verfügbar

2) Szenario B: ADAMTS13 Aktivität *nicht* verfügbar

3) Szenario C: ADAMTS13 Aktivität *verzögert* verfügbar

Diagnostische Strategie für Patienten mit sehr hoher ($\geq 90\%$)
Prä-Test-Wahrscheinlichkeit für eine TTP

Diagnostische Strategie für Patienten mit intermediater oder kleiner Prä-Test-Wahrscheinlichkeit für eine TTP

Was hat “Der” denn nun über Diagnostik erzählt ?

- Biomarker für Diagnose und Verlauf vorhanden (ADAMTS13-Aktivität)
- Schnellteste sind verfügbar, Limitationen
- Zeit bis zur Verfügbarkeit der ADAMTS13 Aktivität hat Einfluss auf Therapieentscheidungen
- Rascher unkomplizierter Kontakt zu einem TTP-Zentrum oder TTP-“Experten” für ambulante Kollegen fulminant wichtig!
- TTP ist eine lebensbedrohliche Erkrankung. Im Zweifelsfall behandeln.

➔ **Prof. Lämmle ist next – WTF ?**

Besten Dank!