

Investigator Initiated Trial

Beta Blockers in Patients with Intermittent Claudication and Arterial Hypertension: Results from the Nebivolol or Metoprolol in Arterial Occlusive Disease (NORMA) Trial Espinola-Klein C., Weisser G., Jagodzinski A., Savvidis S., Warnholtz A., Ostad M., Gori T., Munzel T. Hypertension 2011; 58:148-54.

Multizentrische Studien (Auswahl)

Rivaroxaban or Aspirin for Extended Treatment of Venous Thromboembolism

Weitz JI, Lensing AWA, Prins MH, Bauersachs R, Beyer-Westendorf J, Bounameaux H, Brighton TA, Cohen AT, Davidson BL, Decousus H, Freitas MCS, Holberg G, Kakkar AK, Haskell L, van Bellen B, Pap AF, Berkowitz SD, Verhamme P, Wells PS, Prandoni P; EINSTEIN CHOICE Investigators.

N Engl J Med. 2017; 376: 1211-1222.

Ticagrelor versus Clopidogrel in Symptomatic Peripheral Artery Disease.

Hiatt WR, Fowkes FG1, Heizer G, Berger JS, Baumgartner I, Held P, Katona BG, Mahaffey KW, Norgren L, Jones WS, Blomster J, Millegård M, Reist C, Patel MR; EUCLID Trial Steering Committee and Investigators. N Engl J Med. 2017; 376: 32-40.

Fibrinolysis for patients with intermediate-risk pulmonary embolism.

Meyer G, Vicaut E, Danays T, Agnelli G, Becattini C, Beyer-Westendorf J, Bluhmki E, Bouvaist H, Brenner B, Couturaud F, Dellas C, Empen K, Franca A, Galiè N, Geibel A, Goldhaber SZ, Jimenez D, Kozak M, Kupatt C, Kucher N, Lang IM, Lankeit M, Meneveau N, Pacouret G, Palazzini M, Petris A, Pruszczyk P, Rugolotto M, Salvi A, Schellong S, Sebbane M, Sobkowicz B, Stefanovic BS, Thiele H, Torbicki A, Verschuren F, Konstantinides SV; PEITHO Investigators. N Engl J Med. 2014; 370: 1402-11.

Edoxaban versus warfarin for the treatment of symptomatic venous thromboembolism.

Hokusai-VTE Investigators, Büller HR, Décousus H, Grosso MA, Mercuri M, Middeldorp S, Prins MH, Raskob GE, Schellong SM, Schwocho L, Segers A, Shi M, Verhamme P, Wells P.

N Engl J Med. 2013; 369: 1406-15.

Rivaroxaban for thromboprophylaxis in acutely ill medical patients.

Cohen AT, Spiro TE, Büller HR, Haskell L, Hu D, Hull R, Mebazaa A, Merli G, Schellong S, Spyropoulos AC, Tapson V; MAGELLAN Investigators. N Engl J Med. 2013; 368: 513-23.

Oral rivaroxaban for the treatment of symptomatic pulmonary embolism.

EINSTEIN-PE Investigators, Büller HR, Prins MH, Lensin AW, Decousus H, Jacobson BF, Minar E, Chlumsky J, Verhamme P, Wells P, Agnelli G, Cohen A, Berkowitz SD, Bounameaux H, Davidson BL, Misselwitz F, Gallus AS, Raskob GE, Schellong S, Segers A. N Engl J Med. 2012; 366: 1287-97.

Vorapaxar in the secondary prevention of atherothrombotic events.

Morrow DA, Braunwald E, Bonaca MP, Ameriso SF, Dalby AJ, Fish MP, Fox KA, Lipka LJ, Liu X, Nicolau JC, Ophuis AJ, Paolasso E, Scirica BM, Spinar J, Theroux P, Wiviott SD, Strony J, Murphy SA; TRA 2P-TIMI 50 Steering Committee and Investigators. N Engl J Med. 2012; 366: 1404-13. Oral rivaroxaban for symptomatic venous thromboembolism. EINSTEIN Investigators, Bauersachs R, Berkowitz SD, Brenner B, Buller HR, Decousus H, Gallus AS, Lensing AW, Misselwitz F, Prins MH, Raskob GE, Segers A, Verhamme P, Wells P, Agnelli G, Bounameaux H, Cohen A, Davidson BL, Piovella F, Schellong S. N Engl J Med. 2010; 363: 2499-510.